Space Weather Highlights 21 July 1997 - 27 July 1997

Solar activity was very low to low. Region 8065 (N17, L = 277, class/area Dso/150 on 25 July) produced a long-duration C4/SF flare at 25/2035UT during a brief period of growth. A 1600 SFU Tenflare, and Types II and IV sweep bursts accompanied the flare (estimated Type II shock velocity 0900 km/sec). Activity returned to very low levels following the flare.

Solar wind interplanetary magnetic field (IMF) data were received from the WIND spacecraft a few hours per day (IMF data not available 26 - 27 July due to spacecraft location). Solar wind velocity and density data were received from proton monitor data provided by the University of Maryland from their sensor aboard the SOHO spacecraft. Velocities ranged from 510 - 360 km/sec. Densities were mostly in the 03 - 11 p/cc range, but increased early on 24 July peaking near 025 p/cc. Bz varied from minus 05 - plus 05 nT (GSM). Solar sector orientation was mostly away (phi angle near 135 degrees) through 23 July becoming toward (phi angle near 315 degrees) on 24 July. Orientation was not discernible on 25 July.

There were no significant proton enhancements observed at satellite altitudes.

The greater than 2 MeV electron flux at geosynchronous altitudes was at moderate to normal levels.

The geomagnetic field was quiet to unsettled.

Space Weather Forecast 30 July 1997 - 25 August 1997

Solar activity is expected to be very low.

No significant proton enhancements are expected at satellite altitudes.

The greater than 2 MeV electron flux at geosynchronous altitudes is expected to be normal to moderate.

The geomagnetic field is expected to be predominately quiet to unsettled. However, active levels are expected during 30 July due to a favorably positioned coronal hole.



Daily Solar Data

	Radio	Sun	Sunspot	X-ray				Flares				
	Flux	spot	Area	Background	X-	ray Fl	ux		Op	tical		
Date	10.7 cm	No.	(10 ⁻⁶ hemi.)		C	M	X	S	1	2	3	4
21 July	71	0	0	A5.5	0	0	0	0	0	0	0	0
22 July	72	11	10	A1.5	0	0	0	0	0	0	0	0
23 July	76	31	60	A2.6	0	0	0	0	0	0	0	0
24 July	79	44	90	A5.0	0	0	0	3	0	0	0	0
25 July	80	65	200	B1.2	3	0	0	8	0	0	0	0
26 July	77	58	180	B1.3	0	0	0	1	0	0	0	0
<u>27 July</u>	75	53	180	A5.0	0	0	0	3	0	0	0	0

Daily Particle Data

			Daily I ai	nene Dana					
	I	Proton Fluence	;	Electron Fluence					
	(pro	otons/cm ² -day	-sr)	(elec	trons/cm ² -day-sr)				
Date	>1MeV	>10MeV	>100MeV	>.6MeV	>2MeV >41	<u>MeV</u>			
21 July	1.6E + 5	1.8E+4	4.2E+3		2.6E+7				
22 July	2.8E+5	1.8E+4	4.0E+3		2.4E+7				
23 July	5.0E + 5	1.9E + 4	4.3E+3		2.5E+7				
24 July	5.5E+5	1.8E+4	4.3E+3		3.0E+6				
25 July	2.3E+5	1.9E+4	4.7E + 3		5.4E+5				
26 July	3.5E+5	2.4E+4	4.4E+3	7.2E+5					
27 July	2.9E+5	1.9E+4	4.6E+3		1.0E+6				


Daily Geomagnetic Data

			Duny .	Geomagnene Dan	•		
	M	Iiddle Latitude		High Latitude		Estimated	
	F	redericksburg		College		Planetary	
Date	A	K-indices	A	K-indices	A	K-indices	
21 July	7	1-1-2-1-2-3-2-2	0	0-0-0-0-1-0-1-0	5	1-1-3-1-2-2-1	
22 July	7	1-1-1-2-2-3-2-2	1	1-0-0-1-0-1	4	1-0-0-2-2-2-1	
23 July	7	1-2-2-1-2-2-3	1	1-1-0-0-0-0-0-1	5	1-2-1-0-1-1-3	}
24 July	15	2-2-2-3-4-3-3-4	9	1-1-2-1-5-1-1-1	10	2-1-3-3-3-2-3-3	}
25 July	8	2-2-1-2-2-3-2-2	1	2-1-0-0-1-0-0-0	5	2-2-1-0-1-2-2-2	
26 July	8	2-1-1-1-2-2-3-3	0	0-0-0-0-0-1-1	4	1-1-2-0-1-2-2-2	
27 July	7	2-2-2-1-2-2-2	1	0-0-0-2-0-1-0-1	6	2-1-2-2-2-1-2	<u>'</u>

Alerts and Warnings Issued

Date and Time of Issue (UT)	Type of Alert or Warning	Date and Time of Event (UT)
25 Jul 2055	Type II Radio Emission	25 Jul 2024
25 Jul 2109	Type IV Radio Emission	25 Jul 2018
25 Jul 2118	2695MHz Radio Burst	25 Jul 2018
25 Jul 2132	1- 245MHz Radio Burst	25 Jul 2018
26 Jul 0040	245MHz Radio Noise Storm	25 Jul 0639
27 Jul 1624	$A \ge 20$ Watch	29 Jul
27 Jul 1627	$A \ge 20$ Watch	30 Jul





Twenty-seven Day Outlook

	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
30 Jul	72	20	4	13 Aug	72	5	2
31	70	10	3	14	74	8	3
01 Aug	70	8	3	15	74	8	3
02	70	8	3	16	74	5	2
03	70	8	3	17	74	5	2
04	70	8	3	18	74	5	2
05	70	8	3	19	74	5	2
06	70	8	3	20	74	5	2
07	70	5	2	21	72	5	2
08	70	5	2	22	72	5	2
09	70	5	2	23	72	5	2
10	70	5	2	24	72	5	2
11	72	8	3	25	72	5	2
12	72	5	2				



Energetic Event

	Time (UT)	X-ray	Optical Inform	nation	Peak	Sweep Freq
Date	1/2	Integ	Imp Location	Rgn	Radio Flux	Intensity
	Begin Max Max	Class Flux	Brtns Lat CMD	#	245 2695	II IV

No Event Observed

				Flare List			
						ptical	
_		Time		X-ray	Imp /	Location	Rgn
Date	Begin	Max	End	Class.	Brtns	Lat CMD	#
21 July	0240	0400	0413	B1.8			
21 July	0349 1019	1023	1028	B1.8 B1.0			
22 July		res Observ		D 1.0			
23 July	0701	0706	0724	B1.2			
25 July	0701	0733	0724	B1.2 B1.0			
	0811	0733	0737	B1.0 B2.1			
	0954	0958	1002	B2.1 B2.4			
	1105	1109	1118	B1.3			
	1237	1240	1244	B1.5			
	2028	2031	2034	B1.9			
24 July	0245	0249	0257	B1.3 B1.3			
24 July	0335	0338	0343	B1.3 B1.1			
	0606	0612	0545	D1.1	SF	N27W13	8062
	0656	0657	0714	B2.7	SF	N27W13	8062
	1300	1304	1327	B9.1	SF	N17W37	8065
	1844	1852	1903	B5.0	51	1117 1137	0005
	2153	2200	2222	B2.5			
	2335	2341	2348	B2.5			
25 July	0019	0028	0036	B1.9			
23 July	0417	0417	0420	D 1.)	SF	N20W43	8065
	0512	0536	0610	C1.2	SF	N20W44	8065
	B1040	U1040	A1102	B5.2	SF	N23E80	8066
	1201	1204	1211	B2.2	DI.	1123200	0000
	1232	1238	1246	B3.9			
	1326	1331	1337	B3.1			
	1411	1428	1450	B4.7			
	1507	1515	1520	2	SF	N27W33	8062
	1538	1538	1543	B9.1	SF	N17W51	8065
	1744	1749	1752	B3.3		_,_,,,,	
	1824	1830	1834	B1.6			
	1851	1851	1854		SF	N24E68	8066
	1854	1856	1909	C1.1	SF	N17W52	8065
	1934	1941	1948	B4.6		- · · · - · -	
	2011	2028	2115	C4.4	SF	N16W54	8065
	2332	2336	2344	B6.4			



					O	ptical	
		Time		X-ray	Imp /	Location	Rgn
Date	Begin	Max	End	Class.	Brtns	Lat CMD	#
26 July	0016	0016	0019	B4.3	SF	N22W57	8065
20 July	0243			B2.2	31	1122 11 37	8005
		0252	0257				
	0455	0500	0503	B1.4			
	0528	0531	0534	B1.4			
	0742	0749	0753	B1.6			
	1827	1830	1833	B2.0			
	2106	2110	2115	B1.1			
27 July	0022	0026	0029	B1.1			
·	0148	0154	0159	B1.2			
	0209	0215	0222	B8.3			
	0330	0332	0342	B9.5	SF	N32W48	8062
	0427	0439	0447	B2.6			
	0632	0637	0645	B1.4			
	0813	0815	0823	B2.7	SF	N26W56	8062
	1830	1833	1838	B2.2	SF	N24E50	8066
	2049	2055	2059	B1.4			
	2110	2113	2116	B1.0			

Region Summary

Location	1		Characteri					Flares					
	Helio	Area	Extent	Spot	Spot	Mag	X-ray Optical			ptica	cal		
Date (° Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	C I	M X	S	1	2	3	4
Region	ı 8061												
14 Jul S21W02	800	0010	00	AXX	001	A							
15 Jul S21W15	800												
16 Jul S21W28	800												
17 Jul S21W41	800												
18 Jul S21W54	800												
19 Jul S21W67	800												
20 Jul S21W80	800												
21 Jul S21W93	800												
							0	0 0	0	0	0	0	0

Crossed West Limb.

Absolute heliographic longitude: 008



Region Summary – continued.

	Location	1		Sunspot	Characteri	stics		Flares							
		Helio	Area	Extent	Spot	Spot	Mag		X-ray	1		O	ptica	1	_
Date	(° Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4
	R_{o}	gion 80	62												
17 Iu	1 N23E66	260	0000	00	AXX	001	A								
	1 N23E50	261	0000	00	AXX		A								
	1 N23E32 1 N23E39	261	0000	00	АЛЛ	001	А								
	1 N23E39	261													
	1 N23E13	261													
	1 N23E00	261	0000	00		000									
	1 N27W09	256	0030	06	DRO	007	В								
	1 N27W22	256	0030	06	DRO	004	В				2				
	1 N28W37	258	0020	05	BXO	006	В				1				
	1 N28W52	260	0010	02	AXX	003	A				•				
	1 N28W65	260	0010	01	AXX	002	A				2				
27 54	1 1120 11 05	200	0010	01	7 11 11 1	002	11	0	0	0	5	0	0	0	0
Still or	Disk.							Ü	Ü	Ü		Ü	Ü	Ů	
	ite heliograp	hic long	gitude: 26	51											
1105010	ne nenograp	1011	,	,1											
	Region	2063													
20 Iu	1 S25W12	299	0010	03	BXO	002	В								
	1 S25W12 1 S25W25	299	0010	03	БЛО	002	D								
	1 S25W25 1 S25W38	299													
	1 S25W50	299													
	1 S25W51	299													
	1 S25W77	299													
25 Ju	1 525 11 11	2))						0	0	0	0	0	0	0	0
Died o	n Disk.							O	O	O	O	U	U	U	O
	ite heliograp	hic long	gitude: 29	9											
1105010															
22.1		gion 80		00	4 3737	001									
	1 N22E75	186	0010	00	AXX	001	A								
	l N24E61	186	0030	07	CRO	004	В								
	1 N23E47	187	0040	07	CRO	004	В								
	1 N22E33	188	0010	02	BXO	005	В								
	1 N22E20	188	0010	03	BXO	003	В								
27 Ju	1 N22E04	191	0010	01	AXX	002	A	^			^	•	•	_	•
Still or	n Disk.							U	U	U	0	U	U	U	U

Absolute heliographic longitude: 191



Region Summary - continued.

Location	1		Sunspot (Characteri	stics		Flares							
	Helio	Area	Extent	Spot	Spot	Mag		X-ray	/		О	ptica	ıl	
Date (° Lat ° CMD)	Lon	(10 ⁻⁶ hemi)	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4
Re	gion 800	55												
24 Jul N17W43	277	0020	04	BXO	006	В				1				
25 Jul N18W56	277	0150	08	DSO	011	BG	3			5				
26 Jul N17W70	278	0140	08	DAO	009	В				1				
27 Jul N16W82	277	0150	09	DAO	006	В								
							3	0	0	7	0	0	0	0
Still on Disk.														
Absolute heliograp	hic long	itude: 27	7											
	_													
Region 806	6													
25 Jul N22E73	148	0020	04	BXO	003	В				2				
26 Jul N23E59	149	0020	04	BXO	003	В								
27 Jul N24E46	149	0010	05	BXO	003	В				1				
							0	0	0	3	0	0	0	0

Still on Disk.

Absolute heliographic longitude: 149



Recent Solar Indices (preliminary) of the observed monthly mean values


			ot Number				o Flux	Geomagn	
-	Observed		Ratio	Smooth		**Penticton	Smooth	_Planetary	
Month	SWO	RI	RI/SWO	SWO	RI	10.7 cm	Value	Ap	Value
					1995				
July	23.8	14.5	0.61	28.1	17.0	73.9	76.9	08	12.4
August	25.1	14.3	0.57	25.4	15.4	73.8	76.0	09	12.1
September	16.5	11.8	0.72	22.0	13.4	72.0	74.8	13	11.8
October	31.6	21.1	0.67	19.7	12.1	77.9	73.8	16	11.4
November	15.7	09.0	0.57	18.5	11.4	74.2	73.2	09	10.7
December	16.2	10.0	0.62	17.6	10.8	72.6	72.8	09	10.0
					1996				
January	17.6	11.5	0.55	16.8	10.4	74.5	72.4	09	09.8
February	09.1	04.4	0.48	16.2	10.1	71.5	72.2	10	09.8
March	12.1	09.2	0.76	15.4	09.7	72.7	72.1	11	09.9
April	08.5	04.8	0.60	13.6	08.6	69.3	71.6	11	09.7
May	11.8	05.5	0.47	12.9	08.1*	72.1	71.4	07	09.5
June	18.8	11.8	0.63	13.5	08.6*	69.6	71.8	05	09.4
July	13.2	08.2	0.67	13.4	08.5*	71.2	72.0	07	09.3
August	20.5	14.4	0.68	13.1	08.4*	72.4	72.1	09	09.4
September	02.9	01.6	0.62	13.3	08.5*	69.4	72.3*	15	09.3*
October	02.3	01.8	0.78	14.0	09.0*	69.2	72.6*	13	09.1*
November	26.7	18.6*	0.72*	15.4	10.0*	78.7	73.0*	08	09.1*
December	21.1	12.7*	0.60*	12.7	10.6*	77.8	73.3*	07	09.3*
Longrams	00.0	06.5*	0.72*		1997	74.0		09	
January	09.0	00.5*							
February March	11.3		0.67*			73.8 73.5*		11	
March	14.4	∪8.8*	0.61*			73.5*		08*	
April	24.5	15.8*	0.64*			74.5*		10*	
May	28.6	18.5*				74.6*		08*	
June	22.1	13.1*	0.59			72.4*		08*	

^{*}Preliminary estimates.

The lowest smoothed sunspot indices number for Cycle 21, RI = 12.3, occurred September 1986. The highest smoothed sunspot number for Cycle 22, RI=158.5, occurred July 1989.



^{**} From June 1991 onward, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.



Weekly Geosynchronous Satellite Environment Summary

Week Beginning 21 July 1997


Protons plot contains the five minute averaged integral proton flux (protons/cm²-sec-sr) as measured by GOES-9 (W135) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five minute averaged integral electron flux (electrons/ cm²-sec-sr) with energies greater than 2 MeV at GOES-9.

Hp plot contains the five minute averaged magnetic field H component in nanoteslas (nT) as measured by GOES-9. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis. *Kp* plot contains the estimated planetary 3-hour K-index (derived by the USAF 55th Space Weather Squadron) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA. These data are made available through cooperation from the Geological Survey of Canada (GSC) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWO and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. Hparallel is subject to a more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.








Weekly GOES Satellite X-ray and Proton Plots

Proton plot contains the five minute averaged integral proton flux (protons/cm²-sec-sr) as measured by GOES-9 (W135) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.


X-ray plot contains five minute averaged x-ray flux (watts/m²) as measured by GOES 8 and 9 in two wavelength bands, .05 -.4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band





Coronal hole maps from the National Solar Observatory, Kitt Peak, Arizona These maps are reproductions of the coronal hole contours as derived from the Kitt Peak 1083 nm raw data images. These are preliminary data, where solid (and dashed) lines are used to indicate more (and less) confidence in an inference and are printed whenever they are available at SWO.





Coronal hole maps from the National Solar Observatory, Kitt Peak, Arizona
These maps are reproductions of the coronal hole contours as derived from the Kitt Peak 1083 nm raw data images. These are preliminary data, where solid (and dashed) lines are used to indicate more (and less) confidence in an inference and are printed whenever they are available at SWO.

